Your resource...


Sexually Transmitted Diseases is your resource for the most current, comprehensive information available on all aspects of sexually transmitted disease. Its unique, multidisciplinary perspective provides insight into the latest advances in the field.

- Diagnosis and Treatment
- Etiology Pathology
- Epidemiology Microbiology
- Public Health and Behavior

Published bi-monthly, *Sexually Transmitted Diseases* brings you original articles on newly identified syndromes, new investigations into the mechanism of disease, descriptions of clinical experiences and reports on therapeutic trials. Features include notes

from the CDC, NIH and ASHA, case reports and AVDA news.

Subscribe to the resource you can rely on for the latest, most authoritative information on STDs!



No-Risk Guarantee!

If you're not completely satisfied, you may cancel your subscription at any time and receive a refund for all unmailed issues.

Call us today TOLL FREE 1-800-777-2295 (U.S. except AK) or FAX your order 301-824-7390.

Published bi-monthly Annual subscription rates:

Individual \$125.00 (U.S.) - \$135.00 (Other countries) Institution \$173.00 (U.S.) - \$205.00 (Other countries)

Published by

J.B. Lippincott

A Wolters Kluwer Company



Journals Division J. B. Lippincott 227 East Washington Square Philadelphia, PA 19106-3780

for the latest information on Sexually Transm

Sexually Transmitted Diseases

Journal of the American Venereal Disease Association Julius Schachter, PhD, Editor

Abbreviated prescribing information for DIFLUCAN* (fluconazole) $\label{eq:decomposition} % \begin{center} \b$

Abbreviated prescribing information for DIFLUCAN* (fluconazole)
Presentation: Capsules containing 50mg, 150mg or 200mg fluconazole: intravenous infusion containing fluconazole 2mg/ml in 0.9% sodium chloride solution: Powder for Oral Suspension available as two dosage strengths containing either fluconazole 50mg/5ml or 200mg/5ml on reconstitution with 24ml water. Indications and dosage: Systemic candidiasis: 400mg on the first day followed by 200-400mg once daily. Cryptococcosis, including meningitis: 400mg on the first day followed by 200-400mg once daily. Cryptococcosis, including meningitis: 400mg on the first day followed by 200-400mg once daily. Oropharyngeal candidiasis: 50-100mg once daily for 7-14 days or longer in immunocompromised patients. Other mucosal candidia infections: 50-100mg once daily for 14-30 days. Prevention of fungal infections in neutropenic patients following cytotoxic chemotherapy or radiotherapy: 50-100mg once daily whilst patients are predisposed to such infections. Dermal fungal infections: 50mg once daily for up to 6 weeks (usually 2-4 weeks - see data sheet). Vaginal candidiasis: single 150mg dose. Use in the elderly - as above except for those renally impaired - see data sheet. Use in children - not recommended. Administration: DIFLUCAN may be administered either orally or by intravenous infusion at a rate of approximately 5-10ml/min. The dosages for the two routes are equivalent. Contraindications: Hypersensitivity to fluconazole or related azoles, pregnancy and women of childhearing potential unless adequate contraception is employed. Warnings: Lactation: Not recommended. Renal impairment: dosage reduction may be necessary, see data sheet. Drug interactions: Anticoagulants, cyclosporin, oral sulphonylureas, phenytoin, rifampicin or theophylline. Side-effects: Nausea, abdominal discomfort, diarrhoea, flatulence and rarely anaphylaxis. Legal Category: POM. Basic NHS Cost and Package Quantities: DIFLUCAN capsules in calendar packs containing 7 x 50mg (£16.61, Pl. 57/0315); 100ml (



Pfizer

...WITH BARELY A RIPPLE[†]

ORAL/I.V. fluconazole

JENTLE STRENGTH